arXiv:1008.1439v3 [math.FA] 26 Aug 2010

DIRECT AND INVERSE ESTIMATES FOR COMBINATIONS OF BERNSTEIN POLYNOMIALS WITH ENDPOINT SINGULARITIES

WEN-MING LU AND LIN ZHANG

Abstract. We give direct and inverse theorems for the weighted approximation of functions with endpoint singularities by combinations of Bernstein polynomials by the rth Ditzian-Totik modulus of smoothness $\omega_\phi^r(f,t)_w$ where ϕ is an admissible step-weight function.

1. Introduction

The set of all continuous functions, defined on the interval I, is denoted by $C(I)$. For any $f \in C([0,1])$, the corresponding *Bernstein operators* are defined as follows:

$$B_n(f,x) := \sum_{k=0}^{n} f(\frac{k}{n}) p_{n,k}(x),$$

where

$$p_{n,k}(x) := \binom{n}{k} x^k (1-x)^{n-k},\ k = 0,1,2,\dots,n,\ x \in [0,1].$$

Approximation properties of Bernstein operators have been studied very well (see [2], [4], [5]-[9], [14]-[16], for example). In order to approximate the functions with singularities, Della Vecchia et al. [4] and Yu-Zhao [14] introduced some kinds of *modified Bernstein operators*. Throughout the paper, C denotes a positive constant independent of n and x, which may be different in different cases. Ditzian and Totik [5] extended the method of combinations and defined the following combinations of Bernstein operators:

$$B_{n,r}(f,x) := \sum_{i=0}^{r-1} C_i(n) B_{n_i}(f,x),$$

with the conditions:

(a) $n = n_0 < n_1 < \cdots < n_{r-1} \leqslant Cn$,

(b) $\sum_{i=0}^{r-1} |C_i(n)| \leqslant C$,

(c) $\sum_{i=0}^{r-1} C_i(n) = 1$,

(d) $\sum_{i=0}^{r-1} C_i(n) n_i^{-k} = 0$, for $k = 1,\dots,r-1$.

Now, we can define our new combinations of Bernstein operators as follows:

$$B^*_{n,r}(f,x) := B_{n,r}(F_n,x) = \sum_{i=0}^{r-1} C_i(n) B_{n_i}(F_n,x), \tag{1.1}$$

where $C_i(n)$ satisfy the conditions (a)-(d). For the details, it can be referred to [13].

1991 *Mathematics Subject Classification.* Primary 41A10, Secondary 41A17.

Key words and phrases. Bernstein polynomials; Endpoint singularities; Pointwise approximation; Direct and inverse theorems.

2. The main results

Let $\varphi(x)=\sqrt{x(1-x)}$ and let $\phi:[0,1]\longrightarrow R,\ \phi\neq 0$ be an admissible step-weight function of the Ditzian-Totik modulus of smoothness, that is, ϕ satisfies the following conditions:

(I) For every proper subinterval $[a,b]\subseteq[0,1]$ there exists a constant $M_1\equiv M(a,b)>0$ such that $M_1^{-1}\leqslant\phi(x)\leqslant M_1$ for $x\in[a,b]$.

(II) There are two numbers $\beta(0)\geqslant 0$ and $\beta(1)\geqslant 0$ for which

$$\phi(x)\sim\begin{cases} x^{\beta(0)}, & \text{as } x\to 0+,\\ (1-x)^{\beta(1)}, & \text{as } x\to 1-.\end{cases}$$

($X\sim Y$ means $C^{-1}Y\leqslant X\leqslant CY$ for some C).

Combining condition (**I**) and (**II**) on ϕ, we can deduce that

$$M^{-1}\phi_2(x)\leqslant\phi(x)\leqslant M\phi_2(x),\ x\in[0,1],$$

where $\phi_2(x)=x^{\beta(0)}(1-x)^{\beta(1)}$, and M is a positive constant independent of x.
Let

$$w(x)=x^{\alpha}(1-x)^{\beta},\ \alpha,\ \beta\geqslant 0,\ \alpha+\beta>0,\ 0\leqslant x\leqslant 1.$$

and

$$C_w:=\{f\in C((0,1)):\lim_{x\longrightarrow 1}(wf)(x)=\lim_{x\longrightarrow 0}(wf)(x)=0\}.$$

The *norm* in C_w is defined by $\|wf\|_{C_w}:=\|wf\|=\sup\limits_{0\leqslant x\leqslant 1}|(wf)(x)|$. Define

$$W_{\phi}^{r}:=\{f\in C_w: f^{(r-1)}\in A.C.((0,1)),\ \|w\phi^r f^{(r)}\|<\infty\},$$
$$W_{\varphi,\lambda}^{r}:=\{f\in C_w: f^{(r-1)}\in A.C.((0,1)),\ \|w\varphi^{r\lambda} f^{(r)}\|<\infty\}.$$

For $f\in C_w$, define the *weighted modulus of smoothness* by

$$\omega_{\phi}^{r}(f,t)_w:=\sup_{0<h\leqslant t}\{\|w\Delta_{h\phi}^{r}f\|_{[16h^2,1-16h^2]}+\|w\overrightarrow{\Delta}_{h}^{r}f\|_{[0,16h^2]}+\|w\overleftarrow{\Delta}_{h}^{r}f\|_{[1-16h^2,1]}\},$$

where

$$\Delta_{h\phi}^{r}f(x)=\sum_{k=0}^{r}(-1)^k\binom{r}{k}f(x+(\frac{r}{2}-k)h\phi(x)),$$
$$\overrightarrow{\Delta}_{h}^{r}f(x)=\sum_{k=0}^{r}(-1)^k\binom{r}{k}f(x+(r-k)h),$$
$$\overleftarrow{\Delta}_{h}^{r}f(x)=\sum_{k=0}^{r}(-1)^k\binom{r}{k}f(x-kh).$$

Recently Felten showed the following two theorems in [6]:

Theorem A. Let $\varphi(x)=\sqrt{x(1-x)}$ and let $\phi:[0,1]\longrightarrow R,\ \phi\neq 0$ be an admissible step-weight function of the Ditzian-Totik modulus of smoothness([5]) such that ϕ^2 and φ^2/ϕ^2 are concave. Then, for $f\in C[0,1]$ and $0<\alpha<2$, $|B_n(f,x)-f(x)|\leqslant\omega_{\phi}^{2}(f,n^{-1/2}\frac{\varphi(x)}{\phi(x)})$.

Theorem B. Let $\varphi(x)=\sqrt{x(1-x)}$ and let $\phi:[0,1]\longrightarrow R,\ \phi\neq 0$ be an admissible step-weight function of the Ditzian-Totik modulus of smoothness such that ϕ^2 and φ^2/ϕ^2 are concave. Then, for $f\in C[0,1]$ and $0<\alpha<2$, $|B_n(f,x)-f(x)|=O((n^{-1/2}\frac{\varphi(x)}{\phi(x)})^{\alpha})$ implies $\omega_{\phi}^{2}(f,t)=O(t^{\alpha})$.
Our main results are the following:

Theorem 2.1. *For any* $\alpha,\ \beta > 0,\ \min\{\beta(0),\beta(1)\} \geqslant \frac{1}{2},\ f \in C_w$, *we have*

$$|w(x)\phi^r(x)B^{*(r)}_{n,r-1}(f,x)| \leqslant Cn^{\frac{r}{2}}\|wf\|. \tag{2.1}$$

Theorem 2.2. *For any* $\alpha,\ \beta > 0,\ f \in W^r_\phi,\ \min\{\beta(0),\beta(1)\} \geqslant \frac{1}{2}$, *we have*

$$|w(x)\phi^r(x)B^{*(r)}_{n,r-1}(f,x)| \leqslant C\|w\phi^r f^{(r)}\|. \tag{2.2}$$

Theorem 2.3. *For* $f \in C_w,\ \alpha,\ \beta > 0,\ \min\{\beta(0),\beta(1)\} \geqslant \frac{1}{2},\ \alpha_0 \in (0,r)$, *we have*

$$w(x)|f(x) - B^*_{n,r-1}(f,x)| = O((n^{-\frac{1}{2}}\phi^{-1}(x)\delta_n(x))^{\alpha_0}) \Longleftrightarrow \omega^r_\phi(f,t)_w = O(t^{\alpha_0}). \tag{2.3}$$

3. Lemmas

Lemma 3.1. ([15]) *For any non-negative real* u *and* v, *we have*

$$\sum_{k=1}^{n-1}(\frac{k}{n})^{-u}(1-\frac{k}{n})^{-v}p_{n,k}(x) \leqslant Cx^{-u}(1-x)^{-v}. \tag{3.1}$$

Lemma 3.2. ([4]) *If* $\gamma \in R$, *then*

$$\sum_{k=0}^{n}|k-nx|^\gamma p_{n,k}(x) \leqslant Cn^{\frac{\gamma}{2}}\varphi^\gamma(x). \tag{3.2}$$

Lemma 3.3. *For any* $f \in W^r_\phi,\ \alpha,\ \beta > 0$, *we have*

$$\|w\phi^r F^{(r)}_n\| \leqslant C\|w\phi^r f^{(r)}\|. \tag{3.3}$$

Proof. By symmetry, we only prove the above result when $x \in (0, 1/2]$, the others can be done similarly. Obviously, when $x \in (0, 1/n]$, by [5], we have

$$\begin{aligned} |L^{(r)}_r(f,x)| &\leqslant C|\overrightarrow{\Delta}^r_{\frac{1}{r}}f(0)| \leqslant Cn^{-\frac{r}{2}+1}\int_0^{\frac{r}{n}}u^{\frac{r}{2}}|f^{(r)}(u)|du \\ &\leqslant Cn^{-\frac{r}{2}+1}\|w\phi^r f^{(r)}\|\int_0^{\frac{r}{n}}u^{\frac{r}{2}}w^{-1}(u)\phi^{-r}(u)du. \end{aligned}$$

So

$$|w(x)\phi^r(x)F^{(r)}_n(x)| \leqslant C\|w\phi^r f^{(r)}\|.$$

If $x \in [\frac{1}{n}, \frac{2}{n}]$, we have

$$\begin{aligned} |w(x)\phi^r(x)F^{(r)}_n(x)| &\leqslant |w(x)\phi^r(x)f^{(r)}(x)| + |w(x)\phi^r(x)(f(x)-F_n(x))^{(r)}| \\ &:= I_1 + I_2. \end{aligned}$$

For I_2, we have

$$\begin{aligned} f(x) - F_n(x) &= (\psi(nx-1)+1)(f(x) - L_r(f,x)). \\ w(x)\phi^r(x)|(f(x)-F_n(x))^{(r)}| &= w(x)\phi^r(x)\sum_{i=0}^{r}n^i|(f(x)-L_r(f,x))^{(r-i)}|. \end{aligned}$$

By [5], then

$$|(f(x)-L_r(f,x))^{(r-i)}|_{[\frac{1}{n},\frac{2}{n}]} \leqslant C(n^{r-i}\|f-L_r\|_{[\frac{1}{n},\frac{2}{n}]} + n^{-i}\|f^{(r)}\|_{[\frac{1}{n},\frac{2}{n}]}),\ 0<j<r.$$

Now, we estimate

$$I := w(x)\phi^r(x)|f(x) - L_r(x)|. \tag{3.4}$$

By Taylor expansion, we have

$$f(\frac{i}{n}) = \sum_{u=0}^{r-1}\frac{(\frac{i}{n}-x)^u}{u!}f^{(u)}(x) + \frac{1}{(r-1)!}\int_x^{\frac{i}{n}}(\frac{i}{n}-s)^{r-1}f^{(r)}(s)ds, \tag{3.5}$$

It follows from (3.5) and the identity

$$\sum_{i=1}^{r}(\frac{i}{n})^v l_i(x) = Cx^v,\ v = 0,1,\cdots,r.$$

we have

$$\begin{aligned}L_r(f,x) &= \sum_{i=1}^{r}\sum_{u=0}^{r-1}\frac{(\frac{i}{n}-x)^u}{u!}f^{(u)}(x)l_i(x)+\frac{1}{(r-1)!}\sum_{i=1}^{r}l_i(x)\int_x^{\frac{i}{n}}(\frac{i}{n}-s)^{r-1}f^{(r)}(s)ds\\ &= f(x)+C\sum_{u=1}^{r-1}f^{(u)}(x)(\sum_{v=0}^{u}C_u^v(-x)^{u-v}\sum_{i=1}^{r}(\frac{i}{n})^v l_i(x))\\ &+ \frac{1}{(r-1)!}\sum_{i=1}^{r}l_i(x)\int_x^{\frac{i}{n}}(\frac{i}{n}-s)^{r-1}f^{(r)}(s)ds,\end{aligned}$$

which implies that

$$w(x)\phi^r(x)|f(x)-L_r(f,x)|=\frac{1}{(r-1)!}w(x)\phi^r(x)\sum_{i=1}^{r}l_i(x)\int_x^{\frac{i}{n}}(\frac{i}{n}-s)^{r-1}f^{(r)}(s)ds,$$

since $|l_i(x)|\leqslant C$ for $x\in[0,\frac{2}{n}],\ i=1,2,\cdots,r$. It follows from $\frac{|\frac{i}{n}-s|^{r-1}}{w(s)}\leqslant\frac{|\frac{i}{n}-x|^{r-1}}{w(x)}$, s between $\frac{i}{n}$ and x, then

$$\begin{aligned}w(x)\phi^r(x)|f(x)-L_r(f,x)| &\leqslant Cw(x)\phi^r(x)\sum_{i=1}^{r}\int_x^{\frac{i}{n}}(\frac{i}{n}-s)^{r-1}|f^{(r)}(s)|ds\\ &\leqslant C\phi^r(x)\|w\phi^rf^{(r)}\|\sum_{i=1}^{r}\int_x^{\frac{i}{n}}(\frac{i}{n}-s)^{r-1}\phi^{-r}(s)ds\\ &\leqslant \frac{C}{n^r}\|w\phi^rf^{(r)}\|.\end{aligned}$$

Thus $I\leqslant C\|w\phi^rf^{(r)}\|$. So, we get $I_2\leqslant C\|w\phi^rf^{(r)}\|$. Above all, we have

$$|w(x)\phi^r(x)F_n^{(r)}(x)|\leqslant C\|w\phi^rf^{(r)}\|.$$

□

Lemma 3.4. *If* $f\in W_\phi^r,\ \alpha,\ \beta>0,\ \min\{\beta(0),\beta(1)\}\geqslant\frac{1}{2}$, *then*

$$|w(x)(f(x)-L_r(f,x))|_{[0,\frac{2}{n}]}\leqslant C(\frac{\delta_n(x)}{\sqrt{n}\phi(x)})^r\|w\phi^rf^{(r)}\|. \tag{3.6}$$

$$|w(x)(f(x)-R_r(f,x))|_{[1-\frac{2}{n},1]}\leqslant C(\frac{\delta_n(x)}{\sqrt{n}\phi(x)})^r\|w\phi^rf^{(r)}\|. \tag{3.7}$$

Proof. By Taylor expansion, we have

$$f(\frac{i}{n})=\sum_{u=0}^{r-1}\frac{(\frac{i}{n}-x)^u}{u!}f^{(u)}(x)+\frac{1}{(r-1)!}\int_x^{\frac{i}{n}}(\frac{i}{n}-s)^{r-1}f^{(r)}(s)ds, \tag{3.8}$$

It follows from (3.8) and the identities

$$\sum_{i=1}^{r-1}(\frac{i}{n})^v l_i(x)=Cx^v,\ v=0,1,\ldots,r.$$

we have

$$\begin{aligned}L_r(f,x) &= \sum_{i=1}^{r}\sum_{u=0}^{r-1}\frac{(\frac{i}{n}-x)^u}{u!}f^{(u)}(x)l_i(x)+\frac{1}{(r-1)!}\sum_{i=1}^{r}l_i(x)\int_x^{\frac{i}{n}}(\frac{i}{n}-s)^{r-1}f^{(r)}(s)ds\\ &= f(x)+C\sum_{u=1}^{r-1}f^{(u)}(x)(\sum_{v=0}^{u}C_u^v(-x)^{u-v}\sum_{i=1}^{r}(\frac{i}{n})^v l_i(x))\\ &+ \frac{1}{(r-1)!}\sum_{i=1}^{r}l_i(x)\int_x^{\frac{i}{n}}(\frac{i}{n}-s)^{r-1}f^{(r)}(s)ds,\end{aligned}$$

which implies that

$$w(x)|f(x)-L_r(f,x)| = \frac{1}{(r-1)!}w(x)\sum_{i=1}^{r} l_i(x)\int_x^{\frac{i}{n}}(\frac{i}{n}-s)^{r-1}f^{(r)}(s)ds,$$

since $|l_i(x)| \leqslant C$ for $x \in [0, \frac{2}{n}],\ i = 1, 2, \cdots, r$. It follows from $\frac{|\frac{i}{n}-s|^{r-1}}{w(s)} \leqslant \frac{|\frac{i}{n}-x|^{r-1}}{w(x)}$, s between $\frac{i}{n}$ and x, then

$$\begin{aligned} w(x)|f(x)-L_r(f,x)| &\leqslant Cw(x)\sum_{i=1}^{r}\int_x^{\frac{i}{n}}(\frac{i}{n}-s)^{r-1}|f^{(r)}(s)|ds \\ &\leqslant C\frac{\varphi^r(x)}{\phi^r(x)}\|w\phi^r f^{(r)}\|\sum_{i=1}^{r}\int_x^{\frac{i}{n}}(\frac{i}{n}-s)^{r-1}\varphi^{-r}(s)ds \\ &\leqslant C\frac{\delta_n^r(x)}{\phi^r(x)}\|w\phi^r f^{(r)}\|\sum_{i=1}^{r}\int_x^{\frac{i}{n}}(\frac{i}{n}-s)^{r-1}\varphi^{-r}(s)ds \\ &\leqslant C(\frac{\delta_n(x)}{\sqrt{n}\phi(x)})^r\|w\phi^r f^{(r)}\|. \end{aligned}$$

The proof of (3.7) can be done similarly. □

Lemma 3.5. ([13]) *For every* $\alpha,\ \beta > 0$, *we have*

$$\|wB^{*}_{n,r-1}(f)\| \leqslant C\|wf\|. \tag{3.9}$$

Lemma 3.6. ([17]) *Let* $\min\{\beta(0),\beta(1)\} \geqslant \frac{1}{2}$, *then for* $r \in N,\ 0 < t < \frac{1}{8r}$ *and* $\frac{rt}{2} < x < 1-\frac{rt}{2}$, *we have*

$$\int_{-\frac{t}{2}}^{\frac{t}{2}}\cdots\int_{-\frac{t}{2}}^{\frac{t}{2}}\phi^{-r}(x+\sum_{k=1}^{r}u_k)du_1\cdots du_r \leqslant Ct^r\phi^{-r}(x). \tag{3.10}$$

Lemma 3.7. ([10]) *If* $\alpha,\ \beta > 0$, *for any* $f \in C_w$, *we have*

$$\|wB^{*(r)}_{n,r-1}(f)\| \leqslant Cn^r\|wf\|. \tag{3.11}$$

4. PROOF OF THEOREMS

4.1. **Proof of Theorem 2.1.** When $f \in C_w,\ \min\{\beta(0),\beta(1)\} \geqslant \frac{1}{2}$, we discuss it as follows:

Case 1. If $0 \leqslant \varphi(x) \leqslant \frac{1}{\sqrt{n}}$, by (3.11), we have

$$\begin{aligned} |w(x)\phi^r(x)B^{*(r)}_{n,r-1}(f,x)| &= C\varphi^r(x)\cdot\frac{\phi^r(x)}{\varphi^r(x)}|w(x)B^{*(r)}_{n,r-1}(f,x)| \\ &\leqslant Cn^{\frac{r}{2}}\|wf\|. \end{aligned} \tag{4.1}$$

Case 2. If $\varphi(x) > \frac{1}{\sqrt{n}}$, we have

$$\begin{aligned} &|B^{*(r)}_{n,r-1}(f,x)| = |B^{(r)}_{n,r-1}(F_n,x)| \\ \leqslant\ & (\varphi^2(x))^{-r}\sum_{i=0}^{r-2}\sum_{j=0}^{r}Q_j(x,n_i)C_i(n)n_i^j\sum_{k=0}^{n_i}|(x-\frac{k}{n_i})^jF_n(\frac{k}{n_i})|p_{n_i,k}(x), \end{aligned}$$

By [5], we have
$Q_j(x,n_i) = (n_ix(1-x))^{[\frac{r-j}{2}]}$, and $(\varphi^2(x))^{-r}Q_j(x,n_i)n_i^j \leqslant C(n_i/\varphi^2(x))^{\frac{r+j}{2}}$.

So

$$\begin{aligned} &|w(x)\phi^r(x)B^{*(r)}_{n,r-1}(f,x)| \\ \leqslant{}& Cw(x)\phi^r(x)\sum_{i=0}^{r-2}\sum_{j=0}^{r}(\frac{n_i}{\varphi^2(x)})^{\frac{r+j}{2}}\sum_{k=0}^{n_i}|(x-\frac{k}{n_i})^jF_n(\frac{k}{n_i})|p_{n_i,k}(x) \\ \leqslant{}& Cw(x)\phi^r(x)\|wf\|\sum_{i=0}^{r-2}\sum_{j=0}^{r}(\frac{n_i}{\varphi^2(x)})^{\frac{r+j}{2}}\{\sum_{k=0}^{n_i}(x-\frac{k}{n_i})^{2j}\}^{\frac{1}{2}}\cdot \\ &\{\sum_{k=0}^{n_i}w^{-2}(\frac{k}{n_i})p_{n_i,k}(x)\}^{\frac{1}{2}} \\ \leqslant{}& Cn^{\frac{r}{2}}\|wf\|. \end{aligned} \tag{4.2}$$

It follows from combining with (4.1) and (4.2) that the theorem is proved. □

4.2. **Proof of Theorem 2.2.** When $f\in W^r_\phi$, by [5], we have

$$B^{(r)}_{n,r-1}(F_n,x)=\sum_{i=0}^{r-2}C_i(n)n_i^r\sum_{k=0}^{n_i-r}\overrightarrow{\Delta}^r_{\frac{1}{n_i}}F_n(\frac{k}{n_i})p_{n_i-r,k}(x). \tag{4.3}$$

If $0<k<n_i-r$, we have

$$|\overrightarrow{\Delta}^r_{\frac{1}{n_i}}F_n(\frac{k}{n_i})|\leqslant Cn_i^{-r+1}\int_0^{\frac{r}{n_i}}|F_n^{(r)}(\frac{k}{n_i}+u)|du, \tag{4.4}$$

If $k=0$, we have

$$|\overrightarrow{\Delta}^r_{\frac{1}{n_i}}F_n(0)|\leqslant C\int_0^{\frac{r}{n_i}}u^{r-1}|F_n^{(r)}(u)|du, \tag{4.5}$$

Similarly

$$|\overrightarrow{\Delta}^r_{\frac{1}{n_i}}F_n(\frac{n_i-r}{n_i})|\leqslant Cn_i^{-r+1}\int_{1-\frac{r}{n_i}}^{1}(1-u)^{\frac{r}{2}}|F_n^{(r)}(u)|du. \tag{4.6}$$

By (4.3)-(4.6), we have

$$\begin{aligned} &|w(x)\phi^r(x)B^{*(r)}_{n,r-1}(f,x)| \\ \leqslant{}& Cw(x)\phi^r(x)\|w\phi^rF_n^{(r)}\|\sum_{i=0}^{r-2}\sum_{k=0}^{n_i-r}(w\phi^r)^{-1}(\frac{k^*}{n_i})p_{n_i-r,k}(x), \end{aligned} \tag{4.7}$$

where $k^*=1$ for $k=0$, $k^*=n_i-r-1$ for $k=n_i-r$ and $k^*=k$ for $1<k<n_i-r$. By (3.1), we have

$$\sum_{k=0}^{n_i-r}(w\phi^r)^{-1}(\frac{k^*}{n_i})p_{n_i-r,k}(x)\leqslant C(w\phi^r)^{-1}(x).$$

which combining with (4.7) give

$$|w(x)\phi^r(x)B^{*(r)}_{n,r-1}(f,x)|\leqslant C\|w\phi^rf^{(r)}\|.\Box$$

Combining with the theorem 2.1 and theorem 2.2, we can obtain

Corollary *For any* $\alpha,\ \beta>0,\ 0\leqslant\lambda\leqslant1,$ *we have*

$$|w(x)\varphi^{r\lambda}(x)B^{*(r)}_{n,r-1}(f,x)|\leqslant\begin{cases} Cn^{r/2}\{max\{n^{r(1-\lambda)/2},\varphi^{r(\lambda-1)}(x)\}\}\|wf\|, & f\in C_w, \\ C\|w\varphi^{r\lambda}f^{(r)}\|, & f\in W^r_{w,\lambda}. \end{cases} \tag{4.8}$$

4.3. **Proof of Theorem 2.3.**

4.3.1. *The direct theorem.* We know

$$F_n(t)=F_n(x)+F_n'(t)(t-x)+\cdots+\frac{1}{(r-1)!}\int_x^t(t-u)^{r-1}F_n^{(r)}(u)du, \tag{4.9}$$

$$B_{n,r-1}((\cdot-x)^k,x)=0,\ k=1,2,\cdots,r-1. \tag{4.10}$$

According to the definition of W_ϕ^r, for any $g\in W_\phi^r$, we have $B^*_{n,r-1}(g,x)=B_{n,r-1}(G_n(g),x)$, and $w(x)|G_n(x)-B_{n,r-1}(G_n,x)|=w(x)|B_{n,r-1}(R_r(G_n,t,x),x)|$, thereof $R_r(G_n,t,x)=\int_x^t(t-u)^{r-1}G_n^{(r)}(u)du$, we have

$$\begin{aligned}w(x)|G_n(x)-B_{n,r-1}(G_n,x)|&\leqslant C\|w\phi^rG_n^{(r)}\|w(x)B_{n,r-1}(\int_x^t\frac{|t-u|^{r-1}}{w(u)\phi^r(u)}du,x)\\&\leqslant C\|w\phi^rG_n^{(r)}\|w(x)(B_{n,r-1}(\int_x^t\frac{|t-u|^{r-1}}{\phi^{2r}(u)}du,x))^{\frac{1}{2}}\cdot\\&\qquad(B_{n,r-1}(\int_x^t\frac{|t-u|^{r-1}}{w^2(u)}du,x))^{\frac{1}{2}}.\end{aligned} \tag{4.11}$$

also

$$\int_x^t\frac{|t-u|^{r-1}}{\phi^{2r}(u)}du\leqslant C\frac{|t-x|^r}{\phi^{2r}(x)},\ \int_x^t\frac{|t-u|^{r-1}}{w^2(u)}du\leqslant\frac{|t-x|^r}{w^2(x)}. \tag{4.12}$$

By (3.2), (3.3) and (4.12), we have

$$\begin{aligned}w(x)|G_n(x)-B_{n,r-1}(G_n,x)|&\leqslant C\|w\phi^rG_n^{(r)}\|\phi^{-r}(x)B_{n,r-1}(|t-x|^r,x)\\&\leqslant Cn^{-\frac{r}{2}}\frac{\varphi^r(x)}{\phi^r(x)}\|w\phi^rG_n^{(r)}\|\\&\leqslant Cn^{-\frac{r}{2}}\frac{\delta_n^r(x)}{\phi^r(x)}\|w\phi^rG_n^{(r)}\|\\&=C(\frac{\delta_n(x)}{\sqrt{n}\phi(x)})^r\|w\phi^rG_n^{(r)}\|.\end{aligned} \tag{4.13}$$

By (3.6), (3.7) and (4.13), when $g\in W_\phi^r$, then

$$\begin{aligned}w(x)|g(x)-B^*_{n,r-1}(g,x)|&\leqslant w(x)|g(x)-G_n(g,x)|+w(x)|G_n(g,x)-B^*_{n,r-1}(g,x)|\\&\leqslant|w(x)(g(x)-L_r(g,x))|_{[0,\frac{2}{n}]}+|w(x)(g(x)-R_r(g,x))|_{[1-\frac{2}{n},1]}\\&\quad+\ C(\frac{\delta_n(x)}{\sqrt{n}\phi(x)})^r\|w\phi^rG_n^{(r)}\|\\&\leqslant C(\frac{\delta_n(x)}{\sqrt{n}\phi(x)})^r\|w\phi^rg^{(r)}\|.\end{aligned} \tag{4.14}$$

For $f\in C_w$, we choose proper $g\in W_\phi^r$, by (3.9) and (4.14), then

$$\begin{aligned}w(x)|f(x)-B^*_{n,r-1}(f,x)|&\leqslant w(x)|f(x)-g(x)|+w(x)|B^*_{n,r-1}(f-g,x)|\\&\quad+w(x)|g(x)-B^*_{n,r-1}(g,x)|\\&\leqslant C(\|w(f-g)\|+(\frac{\delta_n(x)}{\sqrt{n}\phi(x)})^r\|w\phi^rg^{(r)}\|)\\&\leqslant C\omega_\phi^r(f,\frac{\delta_n(x)}{\sqrt{n}\phi(x)})_w.\Box\end{aligned}$$

4.3.2. *The inverse theorem.* We define the weighted main-part modulus for $D=R_+$ by(see [5])

$$\Omega_\phi^r(C,f,t)_w=\sup_{0<h\leqslant t}\|w\Delta_{h\phi}^rf\|_{[Ch^*,\infty]},$$

$$\Omega_\phi^r(1,f,t)_w=\Omega_\phi^r(f,t)_w.$$

The main-part K-functional is given by

$$K_{r,\phi}(f,t^r)_w=\sup_{0<h\leqslant t}\inf_g\{\|w(f-g)\|_{[Ch^*,\infty]}+t^r\|w\phi^rg^{(r)}\|_{[Ch^*,\infty]},\ g^{(r-1)}\in A.C.((Ch^*,\infty))\}.$$

By [5], we have

$$C^{-1}\Omega_\phi^r(f,t)_w \leqslant \omega_\phi^r(f,t)_w \leqslant C\int_0^t \frac{\Omega_\phi^r(f,\tau)_w}{\tau}d\tau, \tag{4.15}$$

$$C^{-1}K_{r,\phi}(f,t^r)_w \leqslant \Omega_\phi^r(f,t)_w \leqslant CK_{r,\phi}(f,t^r)_w. \tag{4.16}$$

Proof. Let $\delta > 0$, by (4.16), we choose proper g so that

$$\|w(f-g)\| \leqslant C\Omega_\phi^r(f,\delta)_w,\ \|w\phi^r g^{(r)}\| \leqslant C\delta^{-r}\Omega_\phi^r(f,\delta)_w. \tag{4.17}$$

For $r \in N,\ 0 < t < \frac{1}{8r}$ and $\frac{rt}{2} < x < 1 - \frac{rt}{2}$, we have

$$\begin{aligned}
|w(x)\Delta_{h\phi}^r f(x)| &\leqslant |w(x)\Delta_{h\phi}^r(f(x) - B_{n,r-1}^*(f,x))| + |w(x)\Delta_{h\phi}^r B_{n,r-1}^*(f-g,x)| \\
&\quad + |w(x)\Delta_{h\phi}^r B_{n,r-1}^*(g,x)| \\
&\leqslant \sum_{j=0}^r C_r^j\Big(n^{-\frac12}\frac{\delta_n(x+(\frac r2 - j)h\phi(x))}{\phi(x+(\frac r2 - j)h\phi(x))}\Big)^{\alpha_0} \\
&\quad + \int_{-\frac{h\phi(x)}{2}}^{\frac{h\phi(x)}{2}}\cdots\int_{-\frac{h\phi(x)}{2}}^{\frac{h\phi(x)}{2}} w(x)B_{n,r-1}^{*(r)}(f-g, x+\sum_{k=1}^r u_k)du_1\cdots du_r \\
&\quad + \int_{-\frac{h\phi(x)}{2}}^{\frac{h\phi(x)}{2}}\cdots\int_{-\frac{h\phi(x)}{2}}^{\frac{h\phi(x)}{2}} w(x)B_{n,r-1}^{*(r)}(g, x+\sum_{k=1}^r u_k)du_1\cdots du_r \\
&:= J_1 + J_2 + J_3.
\end{aligned} \tag{4.18}$$

Obviously

$$J_1 \leqslant C(n^{-\frac12}\phi^{-1}(x)\delta_n(x))^{\alpha_0}. \tag{4.19}$$

By (3.11) and (4.17), we have

$$\begin{aligned}
J_2 &\leqslant Cn^r\|w(f-g)\|\int_{-\frac{h\phi(x)}{2}}^{\frac{h\phi(x)}{2}}\cdots\int_{-\frac{h\phi(x)}{2}}^{\frac{h\phi(x)}{2}} du_1\cdots du_r \\
&\leqslant Cn^r h^r\phi^r(x)\|w(f-g)\| \\
&\leqslant Cn^r h^r\phi^r(x)\Omega_\phi^r(f,\delta)_w.
\end{aligned} \tag{4.20}$$

By the first inequality of (4.8), we let $\lambda = 1$, and (3.10) as well as (4.17), then

$$\begin{aligned}
J_2 &\leqslant Cn^{\frac r2}\|w(f-g)\|\int_{-\frac{h\phi(x)}{2}}^{\frac{h\phi(x)}{2}}\cdots\int_{-\frac{h\phi(x)}{2}}^{\frac{h\phi(x)}{2}} \varphi^{-r}(x+\sum_{k=1}^r u_k)du_1\cdots du_r \\
&\leqslant Cn^{\frac r2}h^r\phi^r(x)\varphi^{-r}(x)\|w(f-g)\| \\
&\leqslant Cn^{\frac r2}h^r\phi^r(x)\varphi^{-r}(x)\Omega_\phi^r(f,\delta)_w.
\end{aligned} \tag{4.21}$$

By the second inequality of (3.10) and (4.17), we have

$$\begin{aligned}
J_3 &\leqslant C\|w\phi^r g^{(r)}\|w(x)\int_{-\frac{h\phi(x)}{2}}^{\frac{h\phi(x)}{2}}\cdots\int_{-\frac{h\phi(x)}{2}}^{\frac{h\phi(x)}{2}} w^{-1}(x+\sum_{k=1}^r u_k)\phi^{-r}(x+\sum_{k=1}^r u_k)du_1\cdots du_r \\
&\leqslant Ch^r\|w\phi^r g^{(r)}\| \\
&\leqslant Ch^r\delta^{-r}\Omega_\phi^r(f,\delta)_w.
\end{aligned} \tag{4.22}$$

Now, by (4.18)-(4.22), there exists $M > 0$ so that

$$\begin{aligned}|w(x)\Delta^r_{h\phi}f(x)| &\leqslant C((n^{-\frac{1}{2}}\frac{\delta_n(x)}{\phi(x)})^{\alpha_0} \\ &+\min\{n^{\frac{r}{2}}\frac{\phi^r(x)}{\varphi^r(x)}, n^r\phi^r(x)\}h^r\Omega^r_\phi(f,\delta)_w + h^r\delta^{-r}\Omega^r_\phi(f,\delta)_w) \\ &\leqslant C((n^{-\frac{1}{2}}\frac{\delta_n(x)}{\phi(x)})^{\alpha_0} \\ &+ h^rM^r(n^{-\frac{1}{2}}\frac{\varphi(x)}{\phi(x)} + n^{-\frac{1}{2}}\frac{n^{-1/2}}{\phi(x)})^{-r}\Omega^r_\phi(f,\delta)_w + h^r\delta^{-r}\Omega^r_\phi(f,\delta)_w) \\ &\leqslant C((n^{-\frac{1}{2}}\frac{\delta_n(x)}{\phi(x)})^{\alpha_0} \\ &+h^rM^r(n^{-\frac{1}{2}}\frac{\delta_n(x)}{\phi(x)})^{-r}\Omega^r_\phi(f,\delta)_w + h^r\delta^{-r}\Omega^r_\phi(f,\delta)_w).\end{aligned}$$

When $n \geqslant 2$, we have

$$n^{-\frac{1}{2}}\delta_n(x) < (n-1)^{-\frac{1}{2}}\delta_{n-1}(x) \leqslant \sqrt{2}n^{-\frac{1}{2}}\delta_n(x),$$

Choosing proper $x,\ \delta,\ n \in N$, so that

$$n^{-\frac{1}{2}}\frac{\delta_n(x)}{\phi(x)} \leqslant \delta < (n-1)^{-\frac{1}{2}}\frac{\delta_{n-1}(x)}{\phi(x)},$$

Therefore

$$|w(x)\Delta^r_{h\phi}f(x)| \leqslant C\{\delta^{\alpha_0} + h^r\delta^{-r}\Omega^r_\phi(f,\delta)_w\}.$$

By Borens-Lorentz lemma in [5], we get

$$\Omega^r_\phi(f,t)_w \leqslant Ct^{\alpha_0}. \tag{4.23}$$

So, by (4.15) and (4.23), we get

$$\omega^r_\phi(f,t)_w \leqslant C\int_0^t \frac{\Omega^r_\phi(f,\tau)_w}{\tau}d\tau = C\int_0^t \tau^{\alpha_0-1}d\tau = Ct^{\alpha_0}.$$

□

References

[1] P. L. Butzer, Linear combinations of Bernstein polynomials, Canad. J. Math. 5 (1953), pp. 559-567.

[2] H. Berens and G. Lorentz, Inverse theorems for Bernstein polynomials, Indiana Univ. Math. J. 21 (1972), pp. 693-708.

[3] Z. Ditzian, A global inverse theorem for combinations of Bernstein polynomials, J. Approx. Theory 26 (1979), pp. 277-292.

[4] D. Della Vechhia, G. Mastroianni and J. Szabados, Weighted approximation of functions with endpoint and inner singularities by Bernstein operators, Acta Math. Hungar. 103 (2004), pp. 19-41.

[5] Z. Ditzian and V. Totik, Moduli of Smoothness, Springer-Verlag, Berlin, New York (1987).

[6] M. Felten, Direct and inverse estimates for Bernstein polynomials, Constr. Approx., 14, 459-468.

[7] S. S. Guo, C. X. Li and X. W. Liu, Pointwise approximation for linear combinations of Bernstein operators, J. Approx. Theory 107 (2000), pp. 109-120.

[8] S. S. Guo, H. Tong and G. Zhang, Pointwise weighted approximation by Bernstein operators, Acta Math. Hungar. 101 (2003), pp. 293-311.

[9] G. G. Lorentz, Bernstein Polynomial, University of Toronto Press, Toronto (1953).

[10] W. M. Lu, L. Zhang and M.Y. Chai, Weighted approximation of functions with endpoint singularities by combinations of Bernstein operatorsarXiv: 1007.5436v2 [math.FA] 2 Aug 2010

[11] L. S. Xie, Pointwise simultaneous approximation by combinations of Bernstein operators, J. Approx. Theory 137 (2005), pp. 1-21.

[12] L. S. Xie, The saturation class for linear combinations of Bernstein operators, Arch. Math. 91 (2008), pp. 86-96.

[13] D. S. Yu, Weighted approximation of functions with singularities by combinations of Bernstein operators, J. Applied Mathematics and Computation. 206(2008),pp.906-918.

[14] D. S. Yu and D. J. Zhao, Approximation of functions with singularities by truncated Bernstein operators, Southeast Bull. Math. 30 (2006), pp. 1178-1189.

[15] D. X. Zhou, Rate of convergence for Bernstein operators with Jacobi weights, Acta Math. Sinica 35 (1992), pp. 331-338.
[16] D. X. Zhou, On smoothness characterized by Bernstein type operators, J. Approx. Theory 81 (1994), pp. 303-315.
[17] J. J. Zhang, Z. B. Xu, Direct and inverse approximation theorems with Jacobi weight for combinations and higer derivatives of Baskakov operators(in Chinese), Journal of systems science and mathematical sciences. 2008 28 (1), pp. 30-39.

Wen-ming Lu
School of Science, Hangzhou Dianzi University, Hangzhou, People's Republic of China
E-mail address: `lu_wenming@163.com`

Lin Zhang
Department of Mathematics, Zhejiang University, Hangzhou, People's Republic of China
E-mail address: `godyalin@163.com;linyz@zju.edu.cn`